

N-DIGIT BENFORD DISTRIBUTED RANDOM VARIABLES

AZAR KHOSRAVANI AND CONSTANTIN RASINARIU

ABSTRACT. The scope of this paper is twofold. First, to emphasize the use of the mod 1 map in exploring the digit distribution of random variables. We show that the well-known base and scale invariance of Benford variables is a consequence of their associated mod 1 density functions being uniformly distributed. Second, to introduce a new concept of the n -digit Benford variable. Such a variable is Benford in the first n digits, but it is not guaranteed to have a logarithmic distribution beyond the n -th digit. We conclude the paper by giving a general construction method for n -digit Benford variables, and provide a concrete example.

1. INTRODUCTION

In 1881 Newcomb [1] noticed that the first digit distribution of numerical data is not uniform but rather logarithmic. He had observed that the pages of logarithm tables were more worn out for smaller digits such as 1 and 2 than for larger ones, and concluded that “[the] law of probability of the occurrence of the numbers is such that all mantissae of their logarithms are equally probable.” He explicitly tabulated the probability of occurrence of the first and second digits. Newcomb’s work was rediscovered in 1938 by Benford [2] who explicitly gave the formula for the probability of a number having the first digit d

$$P(d) = \log \left(1 + \frac{1}{d} \right), \quad d = 1, 2, \dots, 9 \quad (1)$$

where $\log$ is used to represent the base 10 logarithm. He gathered empirical evidence for formula (1) by collecting thousands of numbers from diverse datasets, such as the area of the riverbeds, atomic weights of elements, etc..

Pinkham proved [4] that the only digit distribution that is left invariant under scale change of the underlying distribution is the Benford distribution. Scale invariance means that a collection of data has the same digit distribution when multiplied by a constant. In his seminal paper [3], Hill showed that the appropriate domain for the significant digit probability is the smallest collection of positive real subsets that contains all the infinite sets of the form $\bigcup_{n=-\infty}^{\infty} [a, b) \cdot 10^n$. This set denoted by $\mathcal{A}$, is the smallest sigma algebra generated by $D_1, D_2, \dots$, where D_i is the i -th significant-digit function. $D_1 : \mathbb{R}^+ \rightarrow \{1, \dots, 9\}$ and $D_i : \mathbb{R}^+ \rightarrow \{0, 1, \dots, 9\}$ for $i \neq 1$. For example $D_1(2.718) = 2$ and $D_2(2.718) = 7$. Observe that $D_i^{-1}(d) \in \mathcal{A}$ for all i and d [3]. Within this framework, for a random variable Y , the

Key words and phrases. Benford’s law, random variables, mod 1 map, scale invariance, base invariance.

Benford's first digit distribution law can be stated as

$$P(D_1(Y) = d) = \log(1 + d^{-1}) . \quad (2)$$

In general [6], a random variable is Benford if for all $m \in \mathbb{N}$, all $d_1 \in \{1, \dots, 9\}$ and all $d_j \in \{0, 1, \dots, 9\}$

$$P(D_j(Y) = d_j \text{ for } j = 1, 2, \dots, m) = \log \left(1 + \left(\sum_{j=1}^m 10^{m-j} d_j \right)^{-1} \right) . \quad (3)$$

For example, the probability of having digits 8, 4, and 7 as the first, second, and third significant digits respectively, is

$$P(8, 4, 7) = \log \left(1 + \frac{1}{847} \right) \simeq 0.000512 . \quad (4)$$

Hill defined base invariance and proved that base invariance as well as scale invariance, both imply Benford distribution of digits. Leemis et. al [5] investigated several examples of symmetric and non-symmetric distributions that lead to Benford distributed random variables.

This paper is organized as follows. In section 2 we use the mod 1 map to show that the well-known base and scale invariance of Benford distributed random variables is a consequence of $g^\dagger = 1$, where $g^\dagger$ is the associated mod 1 density function. In section 3 we introduce the concept of n -digit Benford distributed random variables which are guaranteed to obey the log distribution in their first n digits, and give a general construction method for such variables. Unless otherwise specified, throughout this paper we assume that the base is 10.

2. THE MOD 1 MAP

Any positive real number y can be written as $y = m \times 10^k$ for some $k \in \mathbb{Z}$, where $1 \leq m < 10$. Then the first digits of y and m are the same: $D_1(y) = D_1(m)$. Let's assume that y is given by a random variable Y with the density function f . Let $X = \log Y$ be the random variable with the density function g .

Definition 2.1. For any real function $g : \mathbb{R} \rightarrow \mathbb{R}$, we define $g^\dagger = g \pmod{1}$ as

$$g^\dagger(x) = \begin{cases} \sum_{k=-\infty}^{\infty} g(x+k) & \forall x \in [0, 1) \\ 0 & \text{otherwise} \end{cases}$$

Lemma 2.2. *The probability of Y having its first digit d is*

$$P(D_1(Y) = d) = \int_{\log(d)}^{\log(d+1)} g^\dagger(x) dx .$$

Proof. Let us consider the real numbers starting with the digit d , i.e., $D_1(y) = d$. These numbers belong to the set $S = \bigcup_{k=-\infty}^{\infty} [d \times 10^k, (d+1) \times 10^k)$. Now consider the random variable $X = \log Y$ with the density function g . The logarithmic function maps S into

$\bigcup_{k=-\infty}^{\infty} [\log d + k, \log(d+1) + k)$. This set modulo 1 is just $[\log d, \log(d+1))$ as illustrated in figure (1) for the case $d = 2$.




FIGURE 1. The modulo 1 set of logarithms of positive numbers starting with 2.

Consequently

$$\begin{aligned}
 P(D_1(Y) = d) &= \sum_{k=-\infty}^{\infty} P(d \times 10^k \leq Y < (d+1) \times 10^k) \\
 &= \sum_{k=-\infty}^{\infty} P(\log d + k \leq X < \log(d+1) + k) \\
 &= P(\log d \leq X(\bmod 1) < \log(d+1)) \\
 &= \int_{\log d}^{\log(d+1)} g^\dagger(x) dx .
 \end{aligned}$$

□

Clearly if $g^\dagger = 1$, one obtains Benford's law $P(D_1(Y) = d) = \log(1 + 1/d)$. The uniformity of $g^\dagger$ resonates with Newcomb's pioneering observation in 1881 [1] that the “probability of the occurrence of the numbers is such that all mantissae of their logarithms are equally probable.”

Lemma 2.2 can be readily generalized for a sequence of prescribed digits $d_1, d_2, \dots, d_n$ where $d_1 \in \{1, 2, \dots, 9\}$ and $d_i \in \{0, 1, \dots, 9\}$ for $i \geq 1$:

$$P(D_1(Y) = d_1, D_2(Y) = d_2, \dots, D_n(Y) = d_n) = \int_{d_1 + \frac{d_2}{10} + \dots + \frac{d_n}{10^n}}^{d_1 + \frac{d_2}{10} + \dots + \frac{d_n + 1}{10^n}} g^\dagger(x) dx . \quad (5)$$

2.1. Scale Invariance, Base Invariance, and the mod 1 map. Scale invariance means that a collection of data keeps its digit distribution when multiplied by a constant. For example, suppose that the prices of goods are Benford distributed, then scale invariance

implies that these prices remain Benford distributed regardless of the currency in which they are converted.

Lemma 2.3. *Scaling of a random variable Y is equivalent to a translation of $X = \log Y$.*

Proof. Let X be a random variable with the density function g and let X_1 be the random variable generated by the translation of X by t units, i.e.; $X_1 = X + t$ and $g_1(x) = g(x - t)$, where g_1 is the density function of X_1 . Define $Y = 10^X$ and $Y_1 = 10^{X_1}$ and let f and f_1 be their corresponding density functions. In terms of cumulative distribution functions we have

$$F_1(y) = G_1(\log y) = G(\log y - t) = G(\log \frac{y}{10^t}) = F(\frac{y}{10^t}).$$

The converse is immediate. $\square$

Due to modular arithmetics, a translation of g will result in a wrap-around effect in $g^\dagger$. Hence, scaling of Y induces a wrap-around of $g^\dagger$. For example, let $X = \log Y$ be the random variable with the density function $g = \text{Triangle}(0, \frac{3}{2}, 3)$, and let X_1 be its translation by t . The effect of scaling of Y on $g^\dagger$ is shown in figure (2).




FIGURE 2. The wrap-around effect on $g^\dagger$ of a translation by $t=0.65$ of the $g = \text{Triangle}(0, \frac{3}{2}, 3)$.

By lemma (2.2) we observe that $P(D_1(Y) = d) \neq P(D_1(Y_1) = d)$ i.e. the first digit distributions of Y and Y_1 are not the same, indicating that Y is not scale invariant. One can see that only a uniform $g^\dagger$ remains unchanged under the wrap-around effect induced by a scaling of Y .

Theorem 2.4. *Only the random variables characterized by $g^\dagger = 1$ are scale invariant.*

Proof. By lemma (2.3) scaling of Y yields a wrap-around effect of $g^\dagger$. For a Benford distribution, we want the same areas under the $g^\dagger$ and $g_1^\dagger$ over the intervals $[0, \log 2)$, $[\log 2, \log 3)$, $\dots$, $[\log 9, 1]$. Here $g_1^\dagger$ denotes the mod 1 projection of g_1 which is the shifted g . Clearly

a uniform $g^\dagger$ will satisfy this constraint. Any other function when translated, will not simultaneously keep the areas under $g^\dagger$ preserved over all of these intervals. Thus the only $g^\dagger$ left unchanged is the uniform one. $\square$

We are now arriving at two important results first proved by Hill [3] concerning scale and base invariance.

Corollary 2.5. *Scale invariance implies Benford's law.*

Proof. In order to have scale invariance, we must have $g^\dagger = 1$. From lemma (2.2) one obtains Benford's law. $\square$

Base invariance means that for any base b , the probability of having d as the first digit is

$$P_b(D_1(Y) = d) = \log_b(1 + 1/d), \quad d = 1, \dots, b-1.$$

Corollary 2.6. *Scale invariance implies base invariance.*

Proof. Scale invariance demands $g^\dagger = 1$. Then, for an arbitrary base b , integrating over the logarithmic intervals $[\log_b d, \log_b(d+1))$ we get Benford's distribution

$$P(D_1(Y) = d) = \log_b(1 + 1/d), \quad d \in \{1, \dots, b-1\}. \quad (6)$$

$\square$

Using the insight gained by the mod 1 map analysis we can construct an infinite class of Benford distributed random variables with non-compact support. To this end, let us consider X with the following density function

$$g(x) = \sum_{n=1}^{\infty} \frac{1}{2^n} [H(x-n) - H(x-n-1)], \quad (7)$$

as illustrated in figure 3. Here $H(x)$ is the Heaviside step function. Since the series




FIGURE 3. Example of g with non-compact support yielding uniform $g^\dagger$.

$\sum_{n=1}^{\infty} \frac{1}{2^n} = 1$, one obtains $g^\dagger = 1$ and thus, $Y = 10^X$ is Benford distributed. A similar construction can be used for any convergent series.

3. n -DIGIT BENFORD VARIABLES

We define a random variable that has a logarithmic distribution in its first n digits to be an n -digit Benford variable. More precisely, such a variable is Benford in the first n digits, but it is not guaranteed to have a logarithmic distribution beyond the n -th digit.

Definition 3.1. Let $n \in \mathbb{N}$. A random variable is n -digit Benford if for all $d_1 \in \{1, \dots, 9\}$ and all $d_j \in \{0, 1, \dots, 9\}$, $2 \leq j \leq n$

$$P\left(D_1(Y) = d_1, \dots, D_n(Y) = d_n\right) = \log \left(1 + \frac{1}{10^{n-1}d_1 + 10^{n-2}d_2 + \dots + d_n}\right). \quad (8)$$

To show that such variables exist, we use (5) to construct a $g^\dagger$ that satisfies (8) for any $n \in \mathbb{N}$. That is, we search for $g^\dagger$ such that

$$\int_{d_1 + \frac{d_2}{10} + \dots + \frac{d_n}{10^n}}^{d_1 + \frac{d_2}{10} + \dots + \frac{d_n + 1}{10^n}} g^\dagger(x) dx = \log \left(1 + \frac{1}{10^{n-1}d_1 + 10^{n-2}d_2 + \dots + d_n}\right). \quad (9)$$

We proceed by the partition $\{\log 1, \log 2, \dots, \log 10\}$ of the $[0, 1)$ interval. Then, an example of $g^\dagger$ which yields a 1-digit Benford distribution is given by

$$g^\dagger(x) = \begin{cases} \frac{\pi}{2} \sin \left(\pi \frac{x - \log k}{\log(1+1/k)} \right) & \log k < x \leq \log(k+1), \quad k = 1, \dots, 9 \\ 0 & \text{otherwise} \end{cases} \quad (10)$$

as illustrated in figure 4. One can easily check that $\int_{\log(d)}^{\log(d+1)} g^\dagger(x) dx = \log(1 + 1/d)$.




FIGURE 4. Example of $g^\dagger$ yielding a 1-digit Benford random variable.

By theorem (2.4) any density function g whose $g^\dagger$ is non-uniform is not scale invariant. Thus the example of equation (10) while yielding a 1-digit Benford distributed variable, is not scale nor base invariant.

Following this idea, one can generalize the example given in (10) in order to produce a $g^\dagger$ which satisfy (9) with an arbitrary large n . For such a construction let

$$0 = a_0 < a_1 < \dots < a_{m-1} < a_m = 1, \quad m = 10^n$$

be a partition of the interval $[0, 1)$. Let $h_j : [0, 1] \rightarrow [0, \infty)$ be a probability density function. For $x \in [0, 1)$ such that $a_j \leq x < a_{j+1}$, let us define

$$g^\dagger = h_j \left(\frac{x - a_j}{a_{j+1} - a_j} \right) . \quad (11)$$

It is easy to see that for all j ,

$$\int_{a_j}^{a_{j+1}} g^\dagger(x) dx = a_{j+1} - a_j .$$

Therefore we have that $g^\dagger$ behaves as the uniform density ($g(x) = 1$ for all $0 \leq x < 1$) when integrated over intervals of the form $[a_k, a_\ell]$ with $0 \leq k < \ell \leq n$.

4. CONCLUSIONS

In this paper we demonstrated the use of mod 1 map in analyzing the digit distribution of random variables. In particular we have shown that a uniform $g^\dagger$ implies scale and base invariance. We have introduced the concept of n -digit Benford, and gave a concrete example of a 1-digit Benford variable. Furthermore, given any density function, we have shown how to use the mod 1 map to construct an example of a n -digit Benford variable.

ACKNOWLEDGEMENTS

We are grateful to the anonymous referee for the valuable and constructive suggestions that helped improve this work. We also would like to thank Prof. Berger for very useful discussions.

REFERENCES

- [1] Newcomb, S. Note on the frequency of use of the different digits in natural numbers. *American Journal of Mathematics* Vol. 4, No. 1 (1881), pp. 39-40.
- [2] Benford, F. The law of anomalous numbers. *Proceedings of the American Philosophical Society* Vol. 78, No. 4 (1938), pp. 551-572.
- [3] Hill, TP. Base-Invariance Implies Benford's Law. *Proceedings of the American Mathematical Society* Vol. 123, No. 3 (1995), pp. 887-895.
- [4] Pinkham, RS. On the Distribution of First Significant Digits. *Annals of Mathematical Statistics* Vol.32, No. 4 (1961), pp. 1223-1230.
- [5] Leemis, LM, Schmeiser, BW and Evans, DL. Survival Distributions Satisfying Benford's Law. *American Statistician* Vol. 54, No. 4 (2000), pp. 236-241.
- [6] Berger, A and Hill, TP. A basic theory of Benford's Law. *Probability Surveys* Vol. 8 (2011), pp. 1-126.

DEPARTMENT OF SCIENCE AND MATHEMATICS, COLUMBIA COLLEGE CHICAGO, CHICAGO, IL 60605
E-mail address: akhosravani@colum.edu

DEPARTMENT OF SCIENCE AND MATHEMATICS, COLUMBIA COLLEGE CHICAGO, CHICAGO, IL 60605
E-mail address: crasinariu@colum.edu